

Southern Bush Duiker

Tragelaphus scriptus roualeyni

Named for hunter and author Roualeyn Gordon-Cumming, its European "discoverer in" 1850.

DESCRIPTION Shoulder height 28-30 inches (71-76 cm). Weight 100-120 pounds (41-55 kg).

A medium-sized bushbuck, brownish gray in color, without stripes and with fewer spots (sometimes very few) than the Chobe bushbuck.

DISTRIBUTION The Limpopo River drainage in eastern Botswana; the Limpopo and Sabi river drainages in southern and southeastern Zimbabwe; adjacent parts of southern Mozambique; and the Transvaal lowveld and northeastern Zululand in South Africa.



Limpopo Bushbuck

*SCI Record Book of Trophy Animals and the
SCI Trophy Records Committee proudly presents
this certificate to:*

Mechislav Klimovich

for taking a Southern Bush Duiker
with a Rifle measuring 14 6/16

On a hunt in *R.S.A., Limpopo* on 04/08/2015
Measured by Mechislav Klimovich on 11/10/2016.

Entry currently ranks *Number 45, a Gold Medal trophy*

*SCI Record Book of Trophy Animals and the
SCI Trophy Records Committee proudly presents
this certificate to:*

Mechislav Klimovich

for taking a Limpopo Bushbuck

with a Rifle measuring 37 6/8

On a hunt in *R.S.A., Limpopo* on 04/08/2015
Measured by Mechislav Klimovich on 11/10/2016.

Entry currently ranks *Number 102, a Bronze Medal trophy*

Sylvicapra grimmia grimmia

DESCRIPTION Shoulder height 23-25 inches (58-64 cm). Weight 33-46 pounds (15-21 kg).

The southern bush duiker has the largest body and by far the longest horns of any bush duiker. The general color is a pale grayish brown, grizzled with black. The grizzled effect is caused by the alternate yellowish and blackish brown color banding of the individual hairs. The lower parts are colored like the back, except that the chin, the insides of the upper legs, and the underside of the tail are whitish. The face is reddish with a dark brown nose stripe. The ears are long, with narrow pointed tips, and the head is long and narrow.

DISTRIBUTION South Africa, Namibia, Botswana, Zimbabwe, Mozambique, Malawi, Zambia, and Katanga Province in southeastern Congo (K).

TAXONOMIC NOTES Includes seven named races: *burchelli*, *cafra*, *grimmia*, *orbicularis*, *shirensis*, *steinhardtii*, and *transvaalensis*, with *grimmia* Linnaeus, 1758 having priority